

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification<sup>5</sup> : C12N 15/62, C07K 13/00 C12P 21/02</p>	<p>A1</p>	<p>(11) International Publication Number: WO 90/13653 (43) International Publication Date: 15 November 1990 (15.11.90)</p>
<p>(21) International Application Number: PCT/GB90/00650 (22) International Filing Date: 26 April 1990 (26.04.90) (30) Priority data: 8909916.2 29 April 1989 (29.04.89) GB (71) Applicant (for all designated States except US): DELTA BIOTECHNOLOGY LIMITED [GB/GB]; Castle Court, Castle Boulevard, Nottingham NG7 1FD (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): BALLANCE, David, James [GB/GB]; 11 South Road, West Bridgford, Nottingham NG2 7AG (GB). (74) Agent: BASSETT, Richard; Eric Potter &amp; Clarkson, St Mary's Court, St Mary's Gate, Nottingham NG1 1LE (GB).</p>		<p>(81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB, GB (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), SE (European patent), US.  Published With international search report.</p>
<p>(54) Title: FUSION PROTEINS CONTAINING N-TERMINAL FRAGMENTS OF HUMAN SERUM ALBUMIN</p> <p>(57) Abstract</p> <p>A fusion polypeptide comprising, as at least part of the N-terminal portion thereof, an N-terminal portion of HSA or a variant thereof and, as at least part of the C-terminal portion thereof, another polypeptide except that, when the said N-terminal portion of HSA is the 1-n portion where n is 369 to 419 or a variant thereof, then the said polypeptide is one of various specified entities, including the 585 to 1578 portion of human fibronectin or a variant thereof. The HSA-like portion may have additional N-terminal residues, such as secretion leader sequences (signal sequences). The C-terminal portion is preferably the 585-1578 portion of human plasma fibronectin. The N-terminal and C-terminal portions may be cleavable to yield the isolated C-terminal portion, with the N-terminal portion having served to facilitate secretion from the host.</p>		

+ See back of page

BEST AVAILABLE COPY

# DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MC	Monaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
DE	Germany, Federal Republic of	LU	Luxembourg	TG	Togo
DK	Denmark			US	United States of America

Fusion proteins containing N-terminal fragments of human serum albumin

The present invention relates to fusion polypeptides where two individual polypeptides or parts thereof are fused to form a single amino acid chain. Such fusion may arise from the expression of a single continuous coding sequence formed by recombinant DNA techniques.

Fusion polypeptides are known, for example those where a polypeptide which is the ultimately desired product of the process is expressed with an N-terminal "leader sequence" which encourages or allows secretion of the polypeptide from the cell. An example is disclosed in EP-A-116 201 (Chiron).

Human serum albumin (HSA) is a known protein found in the blood. EP-A-147 198 (Delta Biotechnology) discloses its expression in a transformed host, in this case yeast. Our earlier application EP-A-322 094 discloses N-terminal fragments of HSA, namely those consisting of residues 1-n where n is 369 to 419, which have therapeutic utility. The application also mentions the possibility of fusing the C-terminal residue of such molecules to other, unnamed, polypeptides.

One aspect of the present invention provides a fusion polypeptide comprising, as at least part of the N-terminal portion thereof, an N-terminal portion of HSA or a variant thereof and, as at least part of the C-terminal portion thereof, another polypeptide except that, when the said N-terminal portion of HSA is the 1-n portion where n is 369 to 419 or a variant thereof then the said polypeptide is (a) the 585 to 1578 portion of human fibronectin or a variant thereof, (b) the 1 to 368 portion of CD4 or a variant thereof, (c) platelet derived growth factor, or a variant thereof, (d) transforming growth factor, or a variant thereof, (e) the 1-261 portion of mature human plasma fibronectin or a variant thereof, (f) the 278-578 portion of mature human plasma fibronectin or a variant thereof, (g) the 1-272 portion of mature human von Willebrand's Factor or a variant thereof, or (h) alpha-1-antitrypsin or a variant thereof.

The N-terminal portion of HSA is preferably the said 1-n portion, the 1-177 portion (up to and including the cysteine), the 1-200 portion (up to but excluding the cysteine) or a portion intermediate 1-177 and 1-200.

The term "human serum albumin" (HSA) is intended to include (but not necessarily to be restricted to) known or yet-to-be-discovered polymorphic forms of HSA. For example, albumin Naskapi has Lys-372 in place of Glu-372 and pro-albumin Christchurch has an altered pro-sequence. The term "variants" is intended to include (but not necessarily to be restricted to) minor artificial variations in sequence (such as molecules lacking one or a few residues, having conservative substitutions or minor insertions of residues, or having minor variations of amino acid structure). Thus polypeptides which have 80%, preferably 85%, 90%, 95% or 99%, homology with HSA are deemed to be "variants". It is also preferred for such variants to be physiologically equivalent to HSA; that is to say, variants preferably share at least one pharmacological utility with HSA. Furthermore, any putative variant which is to be used pharmacologically should be non-immunogenic in the animal (especially human) being treated.

Conservative substitutions are those where one or more amino acids are substituted for others having similar properties such that one skilled in the art of polypeptide chemistry would expect at least the secondary structure, and preferably the tertiary structure, of the polypeptide to be substantially unchanged. For example, typical such

substitutions include asparagine for glutamine, serine for asparagine and arginine for lysine. Variants may alternatively, or as well, lack up to ten (preferably only one or two) intermediate amino acid residues (ie not at the termini of the said N-terminal portion of HSA) in comparison with the corresponding portion of natural HSA; preferably any such omissions occur in the 100 to 369 portion of the molecule (relative to mature HSA itself) (if present). Similarly, up to ten, but preferably only one or two, amino acids may be added, again in the 100 to 369 portion for preference (if present). The term "physiologically functional equivalents" also encompasses larger molecules comprising the said sequence plus a further sequence at the N-terminal (for example, pro-HSA, pre-pro-HSA and met-HSA).

Clearly, the said "another polypeptide" in the fusion compounds of the invention cannot be the remaining portion of HSA, since otherwise the whole polypeptide would be HSA, which would not then be a "fusion polypeptide".

Even when the HSA-like portion is not the said 1-n portion of HSA, it is preferred for the non-HSA portion to be one of the said (a) to (h) entities.

The 1 to 368 portion of CD4 represents the first four disulphide-linked immunoglobulin-like domains of the human T lymphocyte CD4 protein, the gene for and amino acid sequence of which are disclosed in D. Smith et al (1987) Science 328, 1704-1707. It is used to combat HIV infections.

The sequence of human platelet-derived growth factor (PDGF) is described in Collins et al (1985) Nature 316, 748-750. Similarly, the sequence of transforming growth factors  $\beta$  (TGF- $\beta$ ) is described in Derynck et al (1985) Nature 316, 701-705. These growth factors are useful for wound-healing.

A cDNA sequence for the 1-261 portion of Fn was disclosed in EP-A-207 751 (obtained from plasmid pFH6 with endonuclease PvuII). This portion binds fibrin and can be used to direct fused compounds to blood clots.

A cDNA sequence for the 278-578 portion of Fn, which contains a collagen-binding domain, was disclosed by R.J. Owens and F.E. Baralle in 1986 E.M.B.O.J. 5, 2825-2830. This portion will bind to platelets.

The 1-272 portion of von Willebrand's Factor binds and stabilises factor VIII. The sequence is given in Bontham et al, Nucl. Acids Res. 14, 7125-7127.

Variants of alpha-1-antitrypsin include those disclosed by Rosenberg et al (1984) Nature 312, 77-80. In particular, the present invention includes the Pittsburgh variant (Met<sup>358</sup> is mutated to Arg) and the variant where Pro<sup>357</sup> and Met<sup>358</sup> are mutated to alanine and arginine respectively. These compounds are useful in the treatment of septic shock and lung disorders.

Variants of the non-HSA portion of the polypeptides of the invention include variations as discussed above in relation to the HSA portion, including those with conservative amino acid substitutions, and also homologues from other species.

The fusion polypeptides of the invention may have N-terminal amino acids which extend beyond the portion corresponding to the N-terminal portion of HSA. For example, if the HSA-like portion corresponds to an N-terminal portion of mature HSA, then pre-, pro-, or pre-pro sequences may be added thereto, for example the yeast alpha-factor leader sequence. The fused leader portions of WO 90/01063 may be used. The polypeptide which is



fused to the HSA portion may be a naturally-occurring polypeptide, a fragment thereof or a novel polypeptide, including a fusion polypeptide. For example, in Example 3 below, a fragment of fibronectin is fused to the HSA portion via a 4 amino acid linker.

It has been found that the amino terminal portion of the HSA molecule is so structured as to favour particularly efficient translocation and export of the fusion compounds of the invention in eukaryotic cells.

A second aspect of the invention provides a transformed host having a nucleotide sequence so arranged as to express a fusion polypeptide as described above. By "so arranged", we mean, for example, that the nucleotide sequence is in correct reading frame with an appropriate RNA polymerase binding site and translation start sequence and is under the control of a suitable promoter. The promoter may be homologous with or heterologous to the host. Downstream (3') regulatory sequences may be included if desired, as is known. The host is preferably yeast (for example Saccharomyces spp., e.g. S. cerevisiae; Kluyveromyces spp., e.g. K. lactis; Pichia spp.; or Schizosaccharomyces spp., e.g. S. pombe) but may be any

other suitable host such as E. coli, B. subtilis, Aspergillus spp., mammalian cells, plant cells or insect cells.

A third aspect of the invention provides a process for preparing a fusion polypeptide according to the first aspect of the invention by cultivation of a transformed host according to the second aspect of the invention, followed by separation of the fusion polypeptide in a useful form.

A fourth aspect of the invention provides therapeutic methods of treatment of the human or other animal body comprising administration of such a fusion polypeptide.

In the methods of the invention we are particularly concerned to improve the efficiency of secretion of useful therapeutic human proteins from yeast and have conceived the idea of fusing to amino-terminal portions of HSA those proteins which may ordinarily be only inefficiently secreted. One such protein is a potentially valuable wound-healing polypeptide representing amino acids 585 to 1578 of human fibronectin (referred to herein as Fn 585-1578). As we have described in a separate application (filed simultaneously herewith) this molecule contains cell spreading, chemotactic and chemokinetic activities

useful in healing wounds. The fusion polypeptides of the present invention wherein the C-terminal portion is Fn 585-1578 can be used for wound healing applications as biosynthesised, especially where the hybrid human protein will be topically applied. However, the portion representing amino acids 585 to 1578 of human fibronectin can if desired be recovered from the fusion protein by preceding the first amino acid of the fibronectin portion by amino acids comprising a factor X cleavage site. After isolation of the fusion protein from culture supernatant, the desired molecule is released by factor X cleavage and purified by suitable chromatography (e.g. ion-exchange chromatography). Other sites providing for enzymatic or chemical cleavage can be provided, either by appropriate juxtaposition of the N-terminal and C-terminal portions or by the insertion therebetween of an appropriate linker.

At least some of the fusion polypeptides of the invention, especially those including the said CD4 and vWF fragments, PDGF and  $\alpha_1$ AT, also have an increased half-life in the blood and therefore have advantages and therapeutic utilities themselves, namely the therapeutic utility of the non-HSA portion of the molecule. In the case of  $\alpha_1$ AT and others, the compound will normally be administered as

a one-off dose or only a few doses over a short period, rather than over a long period, and therefore the compounds are less likely to cause an immune response.

#### EXAMPLES : SUMMARY

Standard recombinant DNA procedures were as described by Maniatis et al (1982 and recent 2nd edition) unless otherwise stated. Construction and analysis of phage M13 recombinant clones was as described by Messing (1983) and Sanger et al (1977).

DNA sequences encoding portions of human serum albumin used in the construction of the following molecules are derived from the plasmids mHOB12 and pDBD2 (EP-A-322 094, Delta Biotechnology Ltd, relevant portions of which are reproduced below) or by synthesis of oligonucleotides equivalent to parts of this sequence. DNA sequences encoding portions of human fibronectin are derived from the plasmid pFHDEL1, or by synthesis of oligonucleotides equivalent to parts of this sequence. Plasmid pFHDEL1, which contains the complete human cDNA encoding plasma fibronectin, was obtained by ligation of DNA derived from plasmids pFH6, 16, 54, 154 and 1 (EP-A-207 751; Delta Biotechnology Ltd).

This DNA represents an mRNA variant which does not contain the 'ED' sequence and had an 89-amino acid variant of the III-CS region (R.J. Owens, A.R. Kornblihtt and F.E. Baralle (1986) Oxford Surveys on Eukaryotic Genes 3 141-160). The map of this vector is disclosed in Fig. 11 and the protein sequence of the mature polypeptide produced by expression of this cDNA is shown in Fig. 5.

Oligonucleotides were synthesised on an Applied Biosystems 380B oligonucleotide synthesiser according to the manufacturer's recommendations (Applied Biosystems, Warrington, Cheshire, UK).

An expression vector was constructed in which DNA encoding the HSA secretion signal and mature HSA up to and including the 387th amino acid, leucine, fused in frame to DNA encoding a segment of human fibronectin representing amino acids 585 to 1578 inclusive, was placed downstream of the hybrid promoter of EP-A-258 067 (Delta Biotechnology), which is a highly efficient galactose-inducible promoter functional in Saccharomyces cerevisiae. The codon for the 1578th amino acid of human fibronectin was directly followed by a stop codon (TAA) and then the S. cerevisiae phosphoglycerate kinase (PGK) gene transcription terminator. This vector was then introduced into S. cerevisiae by transformation, wherein it directed

the expression and secretion from the cells of a hybrid molecule representing the N-terminal 387 amino acids of HSA C-terminally fused to amino acids 585 to 1578 of human fibronectin.

In a second example a similar vector is constructed so as to enable secretion by S. cerevisiae of a hybrid molecule representing the N-terminal 195 amino acids of HSA C-terminally fused to amino acids 585 to 1578 of human fibronectin.

Aspects of the present invention will now be described by way of example and with reference to the accompanying drawings, in which:

Figure 1 (on two sheets) depicts the amino acid sequence currently thought to be the most representative of natural HSA, with (boxed) the alternative C-termini of HSA(1-n);

Figure 2 (on two sheets) depicts the DNA sequence coding for mature HSA, wherein the sequence included in Linker 3 is underlined;

Figure 3 illustrates, diagrammatically, the construction of mHOB16;

Figure 4 illustrates, diagrammatically, the construction of pHOB31;

Figure 5 (on 6 sheets) illustrates the mature protein sequence encoded by the Fn plasmid pFHDEL1;

Figure 6 illustrates Linker 5, showing the eight constituent oligonucleotides;

Figure 7 shows schematically the construction of plasmid pDBDF2;

Figure 8 shows schematically the construction of plasmid pDBDF5;

Figure 9 shows schematically the construction of plasmid pDBDF9;

Figure 10 shows schematically the construction of plasmid DBDF12, using plasmid pFHDEL1; and

Figure 11 shows a map of plasmid pFHDEL1.

EXAMPLE 1 : HSA 1-387 FUSED TO Fn 585-1578

The following is an account of a preparation of plasmids comprising sequences encoding a portion of HSA, as is disclosed in EP-A-322 094.

The human serum albumin coding sequence used in the construction of the following molecules is derived from the plasmid M13mpl9.7 (EP-A-201 239, Delta Biotechnology Ltd.) or by synthesis of oligonucleotides equivalent to parts of this sequence. Oligonucleotides were synthesised using phosphoramidite chemistry on an Applied Biosystems 380B oligonucleotide synthesizer according to the manufacturer's recommendations (AB Inc., Warrington, Cheshire, England).

An oligonucleotide was synthesised (Linker A) which represented a part of the known HSA coding sequence (Figure 2) from the PstI site (1235-1240, Figure 2) to the codon for valine 381 wherein that codon was changed from GTG to GTC:



## Linker 1

	D	P	H	E	C	Y
5'	GAT	CCT	CAT	GAA	TGC	TAT
3' ACGT	CTA	GGA	GTA	CTT	ACG	ATA

1247

A	K	V	F	D	E	F	K
GCC	AAA	GTG	TTC	GAT	GAA	TTT	AAA
CGG	TTT	CAC	AAG	CTA	CTT	AAA	TTT

1267

P	L	V
CTT	GTC	3'
GGA	CAG	5'

Linker 1 was ligated into the vector M13mp19 (Norranders *et al*, 1983) which had been digested with PstI and HincII and the ligation mixture was used to transfect E.coli strain XL1-Blue (Stratagene Cloning Systems, San Diego, CA). Recombinant clones were identified by their failure to evolve a blue colour on medium containing the chromogenic indicator X-gal (5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactoside) in the presence of IPTG (isopropylthio- $\beta$ -galactoside). DNA sequence analysis of template DNA prepared from bacteriophage particles of recombinant clones identified a molecule with the required DNA sequence, designated mHOB12 (Figure 3).

M13mpl9.7 consists of the coding region of mature HSA in M13mpl9 (Norrander et al, 1983) such that the codon for the first amino acid of HSA, GAT, overlaps a unique XhoI site thus:

		Asp	Ala	
5'	C T C G A G A T G C A			3'
3'	G A G C T C T A C G T			5'
	<u>XhoI</u>			

(EP-A-210 239). M13mpl9.7 was digested with XhoI and made flush-ended by S1-nuclease treatment and was then ligated with the following oligonucleotide (Linker 2):

Linker 2

5'	T C T T T T A T C C A A G C T T G G A T A A A A G A	3'
3'	A G A A A A T A G G T T C G A A C C T A T T T T C T	5'
	<u>HindIII</u>	

The ligation mix was then used to transfect E.coli XL1-Blue and template DNA was prepared from several plaques and then analysed by DNA sequencing to identify a clone, pDBD1 (Figure 4), with the correct sequence.

A 1.1 kb HindIII to PstI fragment representing the 5' end of the HSA coding region and one half of the inserted oligonucleotide linker was isolated from pDBD1 by agarose gel electrophoresis. This fragment was then ligated with double stranded mHOB12 previously digested with HindIII and PstI and the ligation mix was then used to transfect E.coli XL1-Blue. Single stranded template DNA was prepared from mature bacteriophage particles of several plaques. The DNA was made double stranded in vitro by extension from annealed sequencing primer with the Klenow fragment of DNA polymerase I in the presence of deoxynucleoside triphosphates. Restriction enzyme analysis of this DNA permitted the identification of a clone with the correct configuration, mHOB15 (Figure 4).

The following oligonucleotide (Linker 3) represents from the codon for the 382nd amino acid of mature HSA (glutamate, GAA) to the codon for lysine 389 which is followed by a stop codon (TAA) and a HindIII site and then a BamHI cohesive end:

Linker 3

E	E	P	Q	N	L	I	K	J			
5'	GAA	GAG	CCT	CAG	AAT	TTA	ATC	AAA	TAA	GCTTG	3'
3'	CTT	CTC	GGA	GTC	TTA	AAT	TAG	TTT	ATT	CGAACCTAG	5'

This was ligated into double stranded mHOB15, previously digested with HincII and BamHI. After ligation, the DNA was digested with HincII to destroy all non-recombinant molecules and then used to transfect E.coli XL1-Blue. Single stranded DNA was prepared from bacteriophage particles of a number of clones and subjected to DNA sequence analysis. One clone having the correct DNA sequence was designated mHOB16 (Figure 4).

A molecule in which the mature HSA coding region was fused to the HSA secretion signal was created by insertion of Linker 4 into BamHI and XhoI digested M13mp19.7 to form pDBD2 (Figure 4).

#### Linker 4

	M	K	W	V	S	F	
5' GATCC	ATG	AAG	TGG	GTA	AGC	TTT	
	G TAC	TTC	ACC	CAT	TCG	AAA	
I	S	L	L	F	L	F	S
ATT	TCC	CTT	CTT	TTT	CTC	TTT	AGC
TAA	AGG	GAA	GAA	AAA	GAG	AAA	TCG

19

S	A	Y	S	R	G	V	F
TCG	GCT	TAT	TCC	AGG	GGT	GTG	TTT
AGC	CGA	ATA	AGG	TCC	CCA	CAC	AAA

R        R  
CG        3'  
GCAGCT 5'

In this linker the codon for the fourth amino acid after the initial methionine, ACC for threonine in the HSA pre-pro leader sequence (Lawn *et al.*, 1981), has been changed to AGC for serine to create a HindIII site.

A sequence of synthetic DNA representing a part of the known HSA coding sequence (Lawn *et al.*, 1981) (amino acids 382 to 387, Fig. 2), fused to part of the known fibronectin coding sequence (Kornblihtt *et al.*, 1985) (amino acids 585 to 640, Fig. 2), was prepared by synthesising six oligonucleotides (Linker 5, Fig. 6). The oligonucleotides 2, 3, 4, 6, 7 and 8 were phosphorylated using T4 polynucleotide kinase and then the oligonucleotides were annealed under standard conditions in pairs, i.e. 1+8, 2+7, 3+6 and 4+5. The annealed oligonucleotides were then mixed together and ligated with mHOB12 which had previously been digested with the restriction enzymes HincII and EcoRI. The ligation

mixture was then used to transfect E.coli XL1-Blue (Stratagene Cloning Systems, San Diego, CA). Single stranded template DNA was then prepared from mature bacteriophage particles derived from several independent plaques and then was analysed by DNA sequencing. A clone in which a linker of the expected sequence had been correctly inserted into the vector was designated pDBDF1 (Fig. 7). This plasmid was then digested with PstI and EcoRI and the approx. 0.24kb fragment was purified and then ligated with the 1.29kb BamHI-PstI fragment of pDBD2 (Fig. 7) and BamHI + EcoRI digested pUC19 (Yanisch-Perron, et al., 1985) to form pDBDF2 (Fig. 7).

A plasmid containing a DNA sequence encoding full length human fibronectin, pFHDEL1, was digested with EcoRI and XhoI and a 0.77kb EcoRI-XhoI fragment (Fig. 8) was isolated and then ligated with EcoRI and SalI digested M13 mp18 (Norranders et al., 1983) to form pDBDF3 (Fig. 8).

The following oligonucleotide linker (Linker 6) was synthesised, representing from the PstI site at 4784-4791 of the fibronectin sequence of EP-A-207 751 to the codon for tyrosine 1578 (Fig. 5) which is followed by a stop codon (TAA), a HindIII site and then a BamHI cohesive end:

Linker 6

G P D Q T E M T I E G L  
 GGT CCA GAT CAA ACA GAA ATG ACT ATT GAA GGC TTG  
 A CGT CCA GGT CTA GTT TGT CTT TAC TGA TAA CTT CCG AAC

Q P T V E Y Stop  
 CAG CCC ACA GTG GAG TAT TAA GCTTG  
 GTC GGG TGT CAC CTC ATA ATT CGAACCTAG

This linker was then ligated with PstI and HindIII digested pDBDF3 to form pDBDF4 (Fig. 8). The following DNA fragments were then ligated together with BglII digested pKV50 (EP-A-258 067) as shown in Fig. 8: 0.68kb EcoRI-BamHI fragment of pDBDF4, 1.5kb BamHI-StuI fragment of pDBDF2 and the 2.2kb StuI-EcoRI fragment of pFHDEL1. The resultant plasmid pDBDF5 (Fig. 8) includes the promoter of EP-A-258 067 to direct the expression of the HSA secretion signal fused to DNA encoding amino acids 1-387 of mature HSA, in turn fused directly and in frame with DNA encoding amino acids 585-1578 of human fibronectin, after which translation would terminate at the stop codon TAA. This is then followed by the S.cerevisiae PGK gene transcription terminator. The

plasmid also contains sequences which permit selection and maintenance in Escherichia coli and S.cerevisiae (EP-A-258 067).

This plasmid was introduced into S.cerevisiae S150-2B (leu2-3 leu2-112 ura3-52 trp1-289 his3- 1) by standard procedures (Beggs, 1978). Transformants were subsequently analysed and found to produce the HSA-fibronectin fusion protein.

EXAMPLE 2 : HSA 1-195 FUSED TO Fn 585-1578

In this second example the first domain of human serum albumin (amino acids 1-195) is fused to amino acids 585-1578 of human fibronectin.

The plasmid pDBD2 was digested with BamHI and BqlII and the 0.79kb fragment was purified and then ligated with BamHI-digested M13mp19 to form pDBDF6 (Fig. 6). The following oligonucleotide:

5'-C C A A A G C T C G A G G A A C T T C G-3'

was used as a mutagenic primer to create a XhoI site in pDBDF6 by in vitro mutagenesis using a kit supplied by Amersham International PLC. This site was created by



changing base number 696 of HSA from a T to a G (Fig. 2). The plasmid thus formed was designated pDBDF7 (Fig. 9). The following linker was then synthesised to represent from this newly created XhoI site to the codon for lysine 195 of HSA (AAA) and then from the codon for isoleucine 585 of fibronectin to the ends of oligonucleotides 1 and 8 shown in Fig. 6.

Linker 7

```

      D   E   L   R   D   E   G   K   A   S   S   A   K
TC GAT GAA CTT CGG GAT GAA GGG AAG GCT TCG TCT GCC AAA
      A CTT GAA GCC CTA CTT CCC TTC CGA AGC AGA CGG TTT

      I   T   E   T   P   S   Q   P   N   S   H
ATC ACT GAG ACT CCG AGT CAG C
      TAG TGA CTC TGA GGC TCA GTC GGG TTG AGG GTG G
```

This linker was ligated with the annealed oligonucleotides shown in Fig. 3, i.e. 2+7, 3+6 and 4+5 together with XhoI and EcoRI digested pDBDF7 to form pDBDF8 (Fig. 9). Note that in order to recreate the original HSA DNA sequence, and hence amino acid sequence, insertion of linker 7 and the other oligonucleotides into pDBDF7 does not recreate the XhoI site.

The 0.83kb BamHI-StuI fragment of pDBDF8 was purified and then was ligated with the 0.68kb EcoRI-BamHI fragment of pDBDF2 and the 2.22kb StuI-EcoRI fragment of pFHDEL1 into BglII-digested pKV50 to form pDBDF9 (Fig. 9). This plasmid is similar to pDBDF5 except that it specifies only residues 1-195 of HSA rather than 1-387 as in pDBDF5.

When introduced into S.cerevisiae S150-2B as above, the plasmid directed the expression and secretion of a hybrid molecule composed of residues 1-195 of HSA fused to residues 585-1578 of fibronectin.

EXAMPLE 3 : HSA 1-387 FUSED TO Fn 585-1578, AS CLEAVABLE MOLECULE

In order to facilitate production of large amounts of residues 585-1578 of fibronectin, a construct was made in which DNA encoding residues 1-387 of HSA was separated from DNA encoding residues 585-1578 of fibronectin by the sequence

I E G R  
ATT GAA GGT AGA  
TAA CTT CCA TCT

which specifies the cleavage recognition site for the blood clotting Factor X. Consequently the purified secreted product can be treated with Factor X and then the fibronectin part of the molecule can be separated from the HSA part.

To do this two oligonucleotides were synthesised and then annealed to form Linker 8.

Linker 8

E	E	P	Q	N	L	I	E	G
GAA	GAG	CCT	CAG	AAT	TTA	ATT	GAA	GGT
CTT	CTC	GGA	GTC	TTA	AAT	TAA	CTT	CCA

R	I	T	E	T	P	S	Q	P
AGA	ATC	ACT	GAG	ACT	CCG	AGT	CAG	C
TCT	TAG	TGA	CTC	TGA	GGC	TCA	GTC	GGG

N     S     H

TTG   AGG   GTG G

This linker was then ligated with the annealed oligonucleotides shown in Fig. 6, i.e. 2+7, 3+6 and 4+5 into HincII and EcoRI digested mHOE12, to form pDBDF10

(Fig. 7). The plasmid was then digested with PstI and EcoRI and the roughly 0.24kb fragment was purified and then ligated with the 1.29kb BamHI-PstI fragment of pDBD2 and BamHI and EcoRI digested pUC19 to form pDBDF11 (Fig. 10).

The 1.5kb BamHI-StuI fragment of pDBDF11 was then ligated with the 0.68kb EcoRI-BamHI fragment of pDBDF4 and the 2.22kb StuI-EcoRI fragment of pFHDEL1 into BglII-digested pKV50 to form pDBDF12 (Fig. 10). This plasmid was then introduced into S.cerevisiae S150-2B. The purified secreted fusion protein was treated with Factor X to liberate the fibronectin fragment representing residues 585-1578 of the native molecule.

REFERENCES

Beggs, J.D. (1978) Nature 275, 104-109

Kornblihtt et al. (1985) EMBO J. 4, 1755-1759

Lawn, R.M. et al. (1981) Nucl. Acid. Res. 9, 6103-6114

Maniatis, T. et al. (1982) Molecular cloning: A laboratory manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York.

Messing, J. (1983) Methods Enzymol. 101, 20-78

Norlander, J. et al. (1983) Gene 26, 101-106

Sanger, F. et al. (1977) Proc. Natl. Acad. Sci. USA 74, 5463-5467

Yanisch-Perron, C. (1985) Gene 33, 103-119

## CLAIMS

1. A fusion polypeptide comprising, as at least part of the N-terminal portion thereof, an N-terminal portion of HSA or a variant thereof and, as at least part of the C-terminal portion thereof, another polypeptide except that, when the said N-terminal portion of HSA is the 1-n portion where n is 369 to 419 or a variant thereof then the said polypeptide is (a) the 585 to 1578 portion of human fibronectin or a variant thereof, (b) the 1 to 368 portion of CD4 or a variant thereof, (c) platelet derived growth factor or a variant thereof, (d) transforming growth factor  $\beta$  or a variant thereof, (e) the 1-261 portion of mature human plasma fibronectin or a variant thereof, (f) the 278-578 portion of mature human plasma fibronectin or a variant thereof, (g) the 1-272 portion of mature human von Willebrand's Factor or a variant thereof, or (h) alpha-1-antitrypsin or a variant thereof.

2. A fusion polypeptide according to Claim 1 additionally comprising at least one N-terminal amino acid extending beyond the portion corresponding to the N-terminal portion of HSA.
3. A fusion polypeptide according to Claim 1 or 2 wherein there is a cleavable region at the junction of the said N-terminal or C-terminal portions.
4. A fusion polypeptide according to any one of the preceding claims wherein the said C-terminal portion is the 585 to 1578 portion of human plasma fibronectin or a variant thereof.
5. A transformed or transfected host having a nucleotide sequence so arranged as to express a fusion polypeptide according to any one of the preceding claims.
6. A process for preparing a fusion polypeptide by cultivation of a host according to Claim 5, followed by separation of the fusion polypeptide in a useful form.
7. A fusion polypeptide according to any one of Claims 1 to 4 for use in therapy.

FIGURE 1

	10	20
Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly Glu Glu Asn Phe Lys		
	30	40
Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu Gln Gln Cys Pro Phe Glu Asp His Val		
	50	60
Lys Leu Val Asn Glu Val Thr Glu Phe Ala Lys Thr Cys Val Ala Asp Glu Ser Ala Glu		
	70	80
Asn Cys Asp Lys Ser Leu His Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr Leu		
	90	100
Arg Glu Thr Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu Pro Glu Arg Asn Glu		
	110	120
Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn Leu Pro Arg Leu Val Arg Pro Glu Val		
	130	140
Asp Val Met Cys Thr Ala Phe His Asp Asn Glu Glu Thr Phe Leu Lys Lys Tyr Leu Tyr		
	150	160
Glu Ile Ala Arg Arg His Pro Tyr Phe Tyr Ala Pro Glu Leu Leu Phe Phe Ala Lys Arg		
	170	180
Tyr Lys Ala Ala Phe Thr Glu Cys Cys Gln Ala Ala Asp Lys Ala Ala Cys Leu Leu Pro		
	190	200
Lys Leu Asp Glu Leu Arg Asp Glu Gly Lys Ala Ser Ser Ala Lys Gln Arg Leu Lys Cys		
	210	220
Ala Ser Leu Gln Lys Phe Gly Glu Arg Ala Phe Lys Ala Trp Ala Val Ala Arg Leu Ser		
	230	240
Gln Arg Phe Pro Lys Ala Glu Phe Ala Glu Val Ser Lys Leu Val Thr Asp Leu Thr Lys		
	250	260
Val His Thr Glu Cys Cys His Gly Asp Leu Leu Glu Cys Ala Asp Asp Arg Ala Asp Leu		
	270	280
Ala Lys Tyr Ile Cys Glu Asn Gln Asp Ser Ile Ser Ser Lys Leu Lys Glu Cys Cys Glu		
	290	300
Lys Pro Leu Leu Glu Lys Ser His Cys Ile Ala Glu Val Glu Asn Asp Glu Met Pro Ala		
	310	320
Asp Leu Pro Ser Leu Ala Ala Asp Phe Val Glu Ser Lys Asp Val Cys Lys Asn Tyr Ala		
	330	340
Glu Ala Lys Asp Val Phe Leu Gly Met Phe Leu Tyr Glu Tyr Ala Arg Arg His Pro Asp		
	350	360
Tyr Ser Val Val Leu Leu Leu Arg Leu Ala Lys Thr Tyr Glu Thr Thr Leu Glu Lys Cys		
	370	380
Cys Ala Ala Ala Asp Pro His Glu Cys Tyr Ala Lys Val Phe Asp Glu Phe Lys Pro Leu		



2/18

FIGURE 1 Cont.

	350	400
Val Glu Glu Pro Gln Asn Leu Ile Lys Gln Asn Cys Glu Leu Phe Gln Gln Leu Gly Glu		
	410	420
Tyr Lys Phe Gln Asn Ala Leu Leu Val Arg Tyr Thr Lys Lys Val Pro Gln Val Ser Thr		
	430	440
Pro Thr Leu Val Glu Val Ser Arg Asn Leu Gly Lys Val Gly Ser Lys Cys Cys Lys His		
	450	460
Pro Glu Ala Lys Arg Met Pro Cys Ala Glu Asp Tyr Leu Ser Val Val Leu Asn Gln Leu		
	470	480
Cys Val Leu His Glu Lys Thr Pro Val Ser Asp Arg Val Thr Lys Cys Cys Thr Glu Ser		
	490	500
Leu Val Asn Arg Arg Pro Cys Phe Ser Ala Leu Glu Val Asp Glu Thr Tyr Val Pro Lys		
	510	520
Glu Phe Asn Ala Glu Thr Phe Thr Phe His Ala Asp Ile Cys Thr Leu Ser Glu Lys Glu		
	530	540
Arg Gln Ile Lys Lys Gln Thr Ala Leu Val Glu Leu Val Lys His Lys Pro Lys Ala Thr		
	550	560
Lys Glu Gln Leu Lys Ala Val Met Asp Asp Phe Ala Ala Phe Val Glu Lys Cys Cys Lys		
	570	580
Ala Asp Asp Lys Glu Thr Cys Phe Ala Glu Glu Gly Lys Lys Leu Val Ala Ala Ser Gln		
Ala Ala Leu Gly Leu		

SUBSTITUTE SHEET

FIGURE 2 DNA sequence coding for mature HSA

```

      10      20      30      40      50      60      70      80
GATGCACACAAGAGTGAAGTTGCTCATCGGTTTAAAGATTTGGGAGAAGAAAATTTCAAAGCCTTGGTGTGATTGCCCTT
D A E K S E V A H R F K D L G E E N F K A L V L I A F

      90     100     110     120     130     140     150     160
TGCTCAGTATCTTCAGCACTGTCCATTTGAAGATCATGTAAAAATTAGTGAATGAAGTAAGTGAATTTGCAAAAACATGTG
A Q Y L Q Q C P F E D H V K L V N E V T E F A K T C

      170     180     190     200     210     220     230     240
TTGCTGATGAGTCAGCTGAAAATTGTGACAAATCACCTTCATACCTTTTGGAGACAAATATGCACAGTTGCAACTCTT
V A D E S A E N C D K S L H T L F G D K L C T V A T L

      250     260     270     280     290     300     310     320
CGTGAACCTATGCTGAAATGGCTGACTGCTGTGCAAAACAAGAACTTGAGAGAAATGAATGCTTTTGCACACAAAGA
R E T Y G E M A D C C A K Q E P E R N E C F L Q H R D

      330     340     350     360     370     380     390     400
TGACAACCCAAACCTCCCCGATTGGTGAGACCAGAGGTTGATGTGATGTGCACTGCTTTTCATGACAATGAAGAGACAT
D N P N L P R L V R P E V D V M C T A F H D N E E T

      410     420     430     440     450     460     470     480
TTTTGAAAAATACTTATATGAAATGCCAGAAGACATCCTTACTTTTATGCCCGGAACTCCTTTTCTTTGCTAAAAGG
P L K K Y L Y E I A R R H P Y F Y A P E L L F F A K R

      490     500     510     520     530     540     550     560
TATAAAGCTGCTTTTACAGAAATGTTGCCAAGCTGCTGATAAAGCTGCCTGCCTGTTGCCAAAGCTCGATGAACTTCGGGA
Y K A A F T E C C Q A A D K A A C L L P K L D E L R D

      570     580     590     600     610     620     630     640
TGAAGGGAAGGCTTCTGCTGCCAAACAGAGACTCAAATGTGCCAGTCTCCAAAAATTTGGAGAAAGAGCTTCAAAGCAT
E G K A S S A K Q R L K C A S L Q K F G E R A F K A

      650     660     670     680     690     700     710     720
GGGCACTGGCTCGCCTGAGCCAGAGATTCCCAAAGCTGAGTTTGCAGAAGTTTCCAAGTTAGTGACAGATCTTACCAAA
W A V A R L S Q R F P K A E F A E V S K L V T D L T K

      730     740     750     760     770     780     790     800
GTCCACACGGAATGCTGCCATGGAGATCTGCTTGAATGTGCTGATGACAGGGCGGACCTTGCCAAATATATCTGTGAAAA
V M T E C C H G D L L E C A D D R A D L A K Y E C E N

      810     820     830     840     850     860     870     880
TCAGGATTCGATCTCCAGTAAACTGAAGGAATGCTGTGAAAAACCTCTGTGTGAAAAATCCCACTGCATTGCCGAAGTGG
Q D S I S S K L K E C C E K P L L E K S H C I A E V

      890     900     910     920     930     940     950     960
AAAAATGATGAGATGCCTGCTGACTTGCCTTCATTAGCTGCTGATTTTGTGAAAGTAAGGATGTTTGCAAAACTATGCT
E N D E M P A D L P S L A A D F V E S K D V C K N Y A

      970     980     990    1000    1010    1020    1030    1040
GAGGCAAAAGGATGCTCTCCTGGGCACTGTTTTGTATGAATATGCAAGAAAGGCATCCTGATTAATCTCTGCTGCTGCT
E A K D V F L G M F L Y E Y A R R H P D Y S V V L L L

```

## FIGURE 2 Cont.

1050 1060 1070 1080 1090 1100 1110 1120  
GAGACTTCCCAAGACATATGAAACCACTCTAGAGAAGTGCTGTGCCGCTGCAGATCCTCATGAATGCTATGCCAAAGTGT  
R L A K T Y E T T L E K C C A A A D P H E C Y A K V

1130 1140 1150 1160 1170 1180 1190 1200  
TCGATGAATTTAAACCTCTTGTGGAAGAGCCTCAGAAATTAATCAAAACAAAAGTGTGAGCTTTTGGAGAGCTTGGAGAG  
F D E F K P L V E E P Q N L I K Q N C E L F E C L G E

1210 1220 1230 1240 1250 1260 1270 1280  
TACAAATCCGAATGCCGTATTAGTTCTTACACCAAGAAAGTACCCCAAGTGTCAACTCCAACTCTGTAGAGGTCTC  
Y K F Q N A L L V R Y T K K V P Q V S T P T L V E V S

1290 1300 1310 1320 1330 1340 1350 1360  
AAGAAACCTAGGAAAAGTGGGCAACAAATGTTGTAACATCCTGAAGCAAAAAGAATGCCCTGTGCAGAGACTATCTAT  
R N L G K V G S K C C K H P E A K R M P C A E D Y L

1370 1380 1390 1400 1410 1420 1430 1440  
CCGTGGTCTGAACCAAGTTATGTGTGTTGCATGAGAAAACGCCAGTAAGTGACAGAGTCACAAAATGCTGCACAGAGTCC  
S V V L N Q L C V L H E K T P V S D R V C K C C T E S

1450 1460 1470 1480 1490 1500 1510 1520  
TTGGTGAACAGCGGACCATGCTTTTCAGCTCTGGAAGTCGATGAAACATACGTTCCCAAAGAGTTTAAAGCTGAAACATT  
L V N R R P C F S A L E V D E T Y V P K E F N A E T F


1530 1540 1550 1560 1570 1580 1590 1600  
CACCTTCCATGCAGATATATGCCACACTTTCTGAGAAGGAGAGACAAATCAAGAAACAAACTGCCACTTGTGAGCTTGTGA  
T F H A D I C T L S E K E R Q I K K Q T A L V E L V

1610 1620 1630 1640 1650 1660 1670 1680  
AACACAAGCCCAAGGCAACAAAAGAGCAACTGAAAGCTGTTATGGATGATTTCCGAGCTTTTGTAGAGAGGTGCTGCAAG  
K H K P K A T K E Q L K A V M D D F A A F V E K C C K

1690 1700 1710 1720 1730 1740 1750 1760  
GCTGACGATAAGGAGACCTGCTTTGCCGAGGAGGGTAAAAAACTTGTGCTGCAAGTCAAGCTGCCCTAGGCTTATAACA  
A D D K E T C F A E E G K K L V A A S Q A A L G L


1770 1780  
TCTACATTTAAAAGCATCTCAG

FIGURE 2 Construction of mHOB16



SUBSTITUTE SHEET

FIGURE 1 Construction of pROB31



SUBSTITUTE SHEET

7/18

Fig. 5A

Gln	Ala	Gln	Gln	Met	Val	Gln	Pro	Gln	Ser	Pro	Val	Ala	Val	Ser	Gln	Ser	Lys	Pro	Gly	20	
Cys	Tyr	Asp	Asn	Gly	Lys	His	Tyr	Gln	Ile	30	Asn	Gln	Gln	Trp	Glu	Arg	Thr	Tyr	Leu	Gly	40
Asn	Val	Leu	Val	Cys	Thr	Cys	Tyr	Gly	Gly	50	Ser	Arg	Gly	Phe	Asn	Cys	Glu	Ser	Lys	Pro	60
Glu	Ala	Glu	Glu	Thr	Cys	Phe	Asp	Lys	Tyr	70	Thr	Gly	Asn	Thr	Tyr	Arg	Val	Gly	Asp	Thr	80
Tyr	Glu	Arg	Pro	Lys	Asp	Ser	Met	Ile	Trp	90	Asp	Cys	Thr	Cys	Ile	Gly	Ala	Gly	Arg	Gly	100
Arg	Ile	Ser	Cys	Thr	Ile	Ala	Asn	Arg	Cys	110	His	Glu	Gly	Gly	Gln	Ser	Tyr	Lys	Ile	Gly	120
Asp	Thr	Trp	Arg	Arg	Pro	His	Glu	Thr	Gly	130	Gly	Tyr	Met	Leu	Glu	Cys	Val	Cys	Leu	Gly	140
Asn	Gly	Lys	Gly	Glu	Trp	Thr	Cys	Lys	Pro	150	Ile	Ala	Glu	Lys	Cys	Phe	Asp	His	Ala	Ala	160
Gly	Thr	Ser	Tyr	Val	Val	Gly	Glu	Thr	Trp	170	Glu	Lys	Pro	Tyr	Gln	Gly	Trp	Met	Met	Val	180
Asp	Cys	Thr	Cys	Leu	Gly	Glu	Gly	Ser	Gly	190	Ile	Thr	Cys	Thr	Ser	Arg	Asn	Arg	Cys	Val	200
Asn	Asp	Gln	Asp	Thr	Arg	Thr	Ser	Tyr	Arg	210	Ile	Gly	Asp	Thr	Trp	Ser	Lys	Lys	Asp	Asn	220
Arg	Gly	Asn	Leu	Leu	Gln	Cys	Ile	Cys	Thr	230	Gly	Asn	Gly	Arg	Gly	Glu	Trp	Lys	Cys	Glu	240
Arg	His	Thr	Ser	Val	Gln	Thr	Thr	Ser	Ser	250	Gly	Ser	Gly	Pro	Phe	Thr	Asp	Val	Arg	Ala	260
Ala	Val	Tyr	Gln	Pro	Gln	Pro	His	Pro	Gln	270	Pro	Pro	Tyr	Gly	His	Cys	Val	Thr	Asp	Asp	280
Ser	Ely	Val	Val	Tyr	Ser	Val	Gly	Met	Gln	290	Trp	Leu	Lys	Thr	Gln	Gly	Asn	Lys	Gln	Met	300
Leu	Cys	Thr	Cys	Leu	Gly	Asn	Gly	Val	Ser	310	Cys	Gln	Glu	Thr	Ala	Val	Thr	Gln	Thr	Tyr	320
Gly	Gly	Asn	Ser	Asn	Gly	Glu	Pro	Cys	Val	330	Leu	Pro	Phe	Thr	Tyr	Asn	Gly	Arg	Thr	Phe	340
Tyr	Ser	Cys	Thr	Thr	Glu	Gly	Arg	Gln	Asp	350	Gly	His	Leu	Trp	Cys	Ser	Thr	Thr	Ser	Asn	360
Tyr	Glu	Gln	Asp	Gln	Lys	Tyr	Ser	Phe	Cys	370	Thr	Asp	His	Thr	Val	Leu	Val	Gln	Thr	Gln	380
Gly	Gly	Asn	Ser	Asn	Ely	Ala	Leu	Cys	His	390	Phe	Pro	Phe	Leu	Tyr	Asn	Asn	His	Asn	Tyr	400
Thr	Asp	Cys	Thr	Ser	Glu	Gly	Arg	Arg	Asp	410	Asn	Met	Lys	Trp	Cys	Gly	Thr	Thr	Gln	Asn	420

SUBSTITUTE SHEET

8/18

Fig. 5B

Tyr Asp Ala Asp Gln Lys Phe Gly Phe Cys 430 Pro Met Ala Ala His Glu Glu Ile Cys 440 Thr  
 Thr Asn Glu Gly Val Met Tyr Arg Ile Gly Asp Gln Trp Asp Lys Gln His Asp Met Gly 460  
 His Met Met Arg Cys Thr Cys Val Gly Asn Gly Arg Gly Glu Trp Thr Cys Tyr Ala Tyr 480  
 Ser Gln Leu Arg Asp Gln Cys Ile Val Asp Asp Ile Thr Tyr Asn Val Asn Asp Thr Phe 500  
 His Lys Arg His Glu Glu Gly His Met Leu Asn Cys Thr Cys Phe Gly Gln Gly Arg 520  
 Arg Trp Lys Cys Asp Pro Val Asp Gln Cys Gln Asp Ser Glu Thr Gly Thr Phe Tyr 540  
 Ile Gly Asp Ser Trp Glu Lys Tyr Val His Gly Val Arg Tyr Gln Cys Tyr Cys Tyr 560  
 Arg Gly Ile Gly Glu Trp His Cys Gln Pro Leu Gln Thr Tyr Pro Ser Ser Ser Gly 580  
 Val Glu Val Phe Ile Thr Glu Thr Pro Ser Gln Pro Asn Ser His Pro Ile Gln Trp Asn 600  
 Ala Pro Gln Pro Ser His Ile Ser Lys Tyr Ile Leu Arg Trp Arg Pro Lys Asn Ser Val 620  
 Gly Arg Trp Lys Glu Ala Thr Ile Pro Gly His Leu Asn Ser Tyr Thr Ile Lys Gly Leu 640  
 Lys Pro Gly Val Val Tyr Glu Gly Glr Leu Ile Ser Ile Gln Gln Tyr Gly His Gln Glu 660  
 Val Thr Arg Phe Asp Phe Thr Thr Thr Ser Thr Ser Thr Pro Val Thr Ser Asn Thr 680  
 Thr Gly Glu Thr Thr Pro Phe Ser Pro Leu Val Ala Thr Ser Glu Ser Val Thr Glu Ile 700  
 Thr Ala Ser Ser Phe Val Val Ser Trp Val Ser Ala Ser Asp Thr Val Ser Gly Phe Arg 720  
 Val Glu Tyr Glu Leu Ser Glu Glu Gly Asp Glu Pro Gln Tyr Leu Asp Leu Pro Ser Thr 740  
 Ala Thr Ser Val Asn Ile Pro Asp Leu Leu Pro Ely Arg Lys Tyr Ile Val Asn Val Tyr 760  
 Gln Ile Ser Glu Asp Gly Glu Gln Ser Leu Ile Leu Ser Thr Ser Gln Thr Thr Ala Pro 780  
 Asp Ala Pro Pro Asp Pro Thr Val Asp Gln Val Asp Asp Thr Ser Ile Val Val Arg Trp 800  
 Ser Arg Pro Glr Ala Pro Ile Thr Gly Tyr Arg Ile Val Tyr Ser Pro Ser Val Glu Gly 820  
 Ser Ser Thr Glu Leu Asn Leu Pro Glu Thr Ala Asn Ser Val Thr Leu Ser Asp Leu Glu 840  
 FNDEL 1

SUBSTITUTION SHEET

9/18

Fig. 5C

Pro Gly Val Gln Tyr Asn Ile Thr Ile Tyr	Ala Val Glu Glu Asn Gln Glu Ser Thr	860 Pro
Val Val Ile Gln Gln Glu Thr Thr Gly	Pro Arg Ser Asp Thr Val Pro Ser Pro	880 Arg
Aso Leu Gln Phe Val Glu Val Thr Asp Val	Lys Val Thr Ile Met Trp Thr Pro Pro	900 Glu
Ser Ala Val Thr Gly Tyr Arg Val Asp Val	Ile Pro Val Asn Leu Pro Gly Glu His	920 Gly
Gln Arg Leu Pro Ile Ser Arg Asn Thr	Phe Ala Glu Val Thr Gly Leu Ser Pro Gly	940 Val
Thr Tyr Tyr Phe Lys Val Phe Ala Val Ser	His Gly Arg Glu Ser Lys Pro Leu Thr	960 Ala
Gln Gln Thr Thr Lys Leu Asp Ala Pro Thr	Asn Leu Gln Phe Val Asn Glu Thr Asp	980 Ser
Thr Val Leu Val Arg Trp Thr Pro Pro Arg	Ala Gln Ile Thr Gly Tyr Arg Leu Thr	1000 Val
Gly Leu Thr Arg Arg Gly Gln Pro Arg Gln	Tyr Asn Val Gly Pro Ser Val Ser Lys	1020 Tyr
Pro Leu Arg Asn Leu Gln Pro Ala Ser Glu	Tyr Thr Val Ser Leu Val Ala Ile Lys	1040 Gly
Asn Gln Glu Ser Pro Lys Ala Thr Gly Val	Phe Thr Thr Leu Gln Pro Gly Ser Ser	1060 Ile
Pro Pro Tyr Asn Thr Glu Val Thr Glu Thr	Thr Ile Val Ile Thr Trp Thr Pro Ala	1080 Pro
Arg Ile Gly Phe Lys Leu Gly Val Arg Pro	Ser Gln Gly Gly Glu Ala Pro Arg Glu	1100 Val
Thr Ser Asp Ser Gly Ser Ile Val Val Ser	Gly Leu Thr Pro Gly Val Gly Tyr Val	1120 Tyr
Thr Ile Gln Val Leu Arg Asp Gly Gln Glu	Arg Asp Ala Pro Ile Val Asn Lys Val	1140 Val
Thr Pro Leu Ser Pro Pro Thr Asn Leu His	Leu Glu Ala Asn Pro Asp Thr Gly Val	1160 Leu
Thr Val Ser Trp Glu Arg Ser Thr Thr Pro	Asp Ile Thr Gly Tyr Arg Ile Thr Thr	1180 Thr
Pro Thr Asn Gly Gln Gln Gly Asn Ser Leu	Glu Glu Val Val His Ala Asp Gln Ser	1200 Ser
Cys Thr Phe Asp Asn Leu Ser Pro Gly Leu	Glu Tyr Asn Val Ser Val Tyr Thr Val	1220 Lys
Asp Asp Lys Glu Ser Val Pro Ile Ser Asp	Thr Ile Ile Pro Ala Val Pro Pro Thr	1240 Thr
Asp Leu Arg Phe Thr Asn Ile Gly Pro Asp	Thr Met Arg Val Thr Trp Ala Pro Pro	1260 Pro

SUBSTITUTION SHEET



10/18

Fig. 5D

Ser Ile Asp Leu Thr Asn Phe Leu Val Arg Tyr Ser Pro Val Lys Asn Glu Glu Asp Val  
 1270 1280  
 Ala Glu Leu Ser Ile Ser Pro Ser Asp Asn Ala Val Val Leu Thr Asn Leu Leu Pro Gly  
 1290 1300  
 Thr Glu Tyr Val Val Ser Val Ser Val Tyr Glu Gln His Glu Ser Thr Pro Leu Arg  
 1310 1320  
 Gly Arg Gln Lys Thr Gly Leu Asp Ser Pro Thr Gly Ile Asp Phe Ser Asp Ile Thr Ala  
 1330 1340  
 Asn Ser Phe Thr Val His Trp Ile Ala Pro Arg Ala Thr Ile Thr Gly Tyr Arg Ile Arg  
 1350 1360  
 His His Pro Glu His Phe Ser Gly Arg Pro Arg Glu Asp Arg Val Pro His Ser Arg Asn  
 1370 1380  
 Ser Ile Thr Leu Thr Asn Leu Thr Pro Gly Thr Glu Tyr Val Val Ser Ile Val Ala Leu  
 1390 1400  
 Asn Gly Arg Glu Glu Ser Pro Leu Leu Ile Gly Gln Gln Ser Thr Val Ser Asp Val Pro  
 1410 1420  
 Arg Asp Leu Glu Val Val Ala Ala Thr Pro Thr Ser Leu Leu Ile Ser Trp Asp Ala Pro  
 1430 1440  
 Ala Val Thr Val Arg Tyr Tyr Arg Ile Thr Tyr Gly Glu Thr Gly Gly Asn Ser Pro Val  
 1450 1460  
 Gln Glu Phe Thr Val Pro Gly Ser Lys Ser Thr Ala Thr Ile Ser Gly Leu Lys Pro Gly  
 1470 1480  
 Val Asp Tyr Thr Ile Thr Val Tyr Ala Val Thr Gly Arg Gly Asp Ser Pro Ala Ser Ser  
 1490 1500  
 Lys Pro Ile Ser Ile Asn Tyr Arg Thr Glu Ile Asp Lys Pro Ser Gln Met Gln Val Thr  
 1510 1520  
 Asp Val Gln Asp Asn Ser Ile Ser Val Lys Trp Leu Pro Ser Ser Pro Val Thr Gly  
 1530 1540  
 Tyr Arg Val Thr Thr Thr Pro Lys Asn Gly Pro Gly Pro Thr Lys Thr Lys Thr Ala Gly  
 1550 1560  
 Pro Asp Gln Thr Glu Met Thr Ile Glu Gly Leu Gln Pro Thr Val Glu Tyr Val Val Ser  
 1570 1580  
 Val Tyr Ala Gln Asn Pro Ser Gly Glu Ser Gln Pro Leu Val Gln Thr Ala Val Thr Thr  
 1590 1600  
 Ile Pro Ala Pro Thr Asp Leu Lys Phe Thr Thr Gln Val Thr Pro Thr Ser Leu Ser Ala Gln  
 1610 1620  
 Trp Thr Pro Pro Asn Val Gln Leu Thr Gly Tyr Arg Val Arg Val Thr Pro Lys Glu Lys  
 1630 1640  
 Thr Gly Pro Met Lys Glu Ile Asn Leu Ala Pro Asp Ser Ser Val Val Val Ser Gly  
 1650 1660  
 Leu Met Val Ala Thr Lys Tyr Glu Val Ser Val Tyr Ala Leu Lys Asp Thr Leu Thr Ser  
 1670 1680  
 FNDEL I

SUBSTITUTE SHEET

11/18

Fig. 5E

Arg Pro Ala Gln Gly Val Thr Thr Leu Glu Asn Val Ser Pro Pro Arg Arg Ala Arg 1700  
 Val Thr Asp Ala Thr Glu Thr Thr Ile Thr Ile Ser Trp Arg Thr Lys Thr Glu Thr Ile 1720  
 Thr Gly Phe Gln Val Asp Ala Val Pro Ala Asn Gly Gln Thr Pro Ile Gln Arg Thr Ile 1740  
 Lys Pro Asp Val Arg Ser Tyr Thr Ile Thr Gly Leu Gln Pro Gly Thr Asp Tyr Lys Ile 1760  
 Tyr Leu Tyr Thr Leu Asn Asp Asn Ala Arg Ser Ser Pro Val Val Ile Asp Ala Ser Thr 1780  
 Ala Ile Asp Ala Pro Ser Asn Lau Arg Phe Lau Ala Thr Thr Pro Asn Ser Leu Leu Val 1800  
 Ser Trp Gln Pro Pro Arg Ala Arg Ile Thr Gly Tyr Ile Ile Lys Tyr Glu Lys Pro Gly 1820  
 Ser Pro Pro Arg Glu Val Val Pro Arg Pro Arg Pro Gly Val Thr Glu Ala Thr Ile Thr 1840  
 Gly Leu Glu Pro Gly Thr Glu Tyr Thr Ile Tyr Val Ile Ala Leu Lys Asn Gln Lys 1860  
 Ser Glu Pro Leu Ile Gly Arg Lys Lys Thr Aso Glu Leu Pro Gln Leu Val Thr Leu Pro 1880  
 His Pro Asn Leu His Gly Pro Glu Ile Leu Asp Val Pro Ser Thr Val Gln Lys Thr 1900  
 Phe Val Thr His Pro Gly Tyr Asp Thr Gly Asn Gly Ile Gln Leu Pro Gly Thr Ser Gly 1920  
 Gln Gln Pro Ser Val Gly Gln Gln Met Ile Phe Glu Glu His Gly phe Arg Arg Thr 1940  
 Pro Pro Thr Thr Ala Thr Pro Ile Arg His Arg Pro Arg Pro Tyr Pro Pro Asn Val Ala 1960  
 Leu Ser Gln Thr Thr Ile Ser Trp Ala Pro Phe Gln Aso Thr Ser Glu Tyr Ile Ile Ser 1980  
 Cys His Pro Val Gly Thr Asp Glu Glu Pro Leu Gln Phe Arg Val Pro Gly Thr Ser 2000  
 Ser Ala Thr Leu Thr Gly Leu Thr Arg Gly Ala Thr Tyr Asn Ile Ile Val Glu Ala Leu 2020  
 Lys Asp Gln Gln Arg His Lys Val Arg Glu Glu Val Val Thr Val Gly Asn Ser Val Asn 2040  
 Glu Gly Leu Asn Gln Pro Thr Asp Asp Ser Cys Phe Asp Pro Tyr Thr Val Ser His Tyr 2060  
 Ala Val Gly Asp Glu Trp Glu Arg Met Ser Glu Ser Gly Phe Lys Leu Leu Cys Gln Cys 2080  
 Leu Ser Phe Gly Ser Gly His Phe Arg Cys Asp Ser Ser Arg Trp Cys His Asp Asn Gly 2100

SUBSTITUTE SHEET

12/18

Val Asn Tyr Lys Ile Gly Glu Lys Trp Asp<sup>2110</sup> Arg Gln Gly Glu Asn Gly Gln Met Met Ser<sup>2120</sup>  
 Cys Thr Cys Leu Gly Asn Gly Lys Gly Glu<sup>2130</sup> Phe Lys Cys Asp Pro His Glu Ala Thr Cys<sup>2140</sup>  
 Tyr Asp Asp Gly Lys Thr Tyr His Val Gly<sup>2150</sup> Glu Gln Trp Gln Lys Glu Tyr Leu Gly Ala<sup>2160</sup>  
 Ile Cys Ser Cys Thr Cys Phe Gly Gly Gln Arg Gly Trp Arg Cys Asp Asn Cys Arg Arg<sup>2180</sup>  
 Pro Gly Gly Glu Pro Ser Pro Glu Gly Thr<sup>2190</sup> Thr Gly Gln Ser Tyr Asn Gln Tyr Ser Gln<sup>2200</sup>  
 Arg Tyr His Gln Arg Thr Asn Thr Asn Val Asn Cys Pro Ile Glu Cys Phe Met Pro Leu<sup>2220</sup>  
 Asp Val Gln Ala Asp Arg Glu Asp Ser Arg Glu<sup>2230</sup>

Fig. 5F

SEQUENCE SHEET

13/18




Figure 6 Linker 5 showing the eight constituent oligonucleotides

SUPPLEMENT SHEET

14/18




Fig. 7 Construction of pDBDF2

SUPPLEMENT SHEET




Fig. 8 Construction of pDBDF5

SUBSTITUTE SHEET




Fig. 9 Construction of pDBDF9

SUPPLEMENTARY SHEET





Fig. 10 Construction of pDBDF12



Figure 1

Name: pFHDEL1  
Vector: pUC18 Amp<sup>r</sup> 2860bp  
Insert: hFNCdNA - 7630bp



SUBSTITUTE SHEET

# INTERNATIONAL SEARCH REPORT

International Application No PCT/GB 90/00650

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) * According to International Patent Classification (IPC) or to both National Classification and IPC IPC <sup>5</sup> : C 12 N 15/62, C 07 K 13/00, C 12 P 21/02											
<b>II. FIELDS SEARCHED</b> Minimum Documentation Searched ? Classification System   Classification Symbols IPC <sup>5</sup>   C 12 N, C 12 P, C 07 K Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched *											
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT</b> <table border="1"> <thead> <tr> <th>Category *</th> <th>Citation of Document, ** with indication, where appropriate, of the relevant passages **</th> <th>Relevant to Claim No. **</th> </tr> </thead> <tbody> <tr> <td>A</td> <td>EP, A, 0308381 (SKANDIGEN et al.) 22 March 1989</td> <td></td> </tr> <tr> <td>T</td> <td>EP, A, 0322094 (DELTA BIOTECHNOLOGY LTD) 28 June 1989 (cited in the application)</td> <td></td> </tr> </tbody> </table>			Category *	Citation of Document, ** with indication, where appropriate, of the relevant passages **	Relevant to Claim No. **	A	EP, A, 0308381 (SKANDIGEN et al.) 22 March 1989		T	EP, A, 0322094 (DELTA BIOTECHNOLOGY LTD) 28 June 1989 (cited in the application)	
Category *	Citation of Document, ** with indication, where appropriate, of the relevant passages **	Relevant to Claim No. **									
A	EP, A, 0308381 (SKANDIGEN et al.) 22 March 1989										
T	EP, A, 0322094 (DELTA BIOTECHNOLOGY LTD) 28 June 1989 (cited in the application)										
<p>* Special categories of cited documents: **</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"Δ" document member of the same patent family</p>											
<b>IV. CERTIFICATION</b> <table border="1"> <tr> <td>Date of the Actual Completion of the International Search</td> <td>Date of Mailing of this International Search Report</td> </tr> <tr> <td>10th July 1990</td> <td>09. 08. 90</td> </tr> <tr> <td>International Searching Authority</td> <td>Signature of Authorized Officer</td> </tr> <tr> <td>EUROPEAN PATENT OFFICE</td> <td>M. SOTELO</td> </tr> </table>			Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	10th July 1990	09. 08. 90	International Searching Authority	Signature of Authorized Officer	EUROPEAN PATENT OFFICE	M. SOTELO	
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report										
10th July 1990	09. 08. 90										
International Searching Authority	Signature of Authorized Officer										
EUROPEAN PATENT OFFICE	M. SOTELO										

**ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO.**

GB 9000650  
SA 36670

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.  
The members are as contained in the European Patent Office EDP file on 31/07/90  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A- 0308381	22-03-89	SE-B- 459586	17-07-89
		AU-A- 2420488	17-04-89
		SE-A- 8703539	15-03-89
		WO-A- 8902467	23-03-89
EP-A- 0322094	28-06-89	AU-A- 2404688	18-05-89

EPO FORM P017

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**